arXiv:0707.3360v1 [math.DG] 23 Jul 2007

# SOME CONSTRUCTIONS OF ALMOST PARA-HYPERHERMITIAN STRUCTURES ON MANIFOLDS AND TANGENT BUNDLES

STERE IANUŞ, GABRIEL EDUARD VÎLCU

Abstract. In this paper we give some examples of almost para-hyperhermitian structures on the tangent bundle of an almost product manifold, on the product manifold $M\times\mathbb{R}$, where $M$ is a manifold endowed with a mixed 3-structure and on the circle bundle over a manifold with a mixed 3-structure.
*AMS Mathematics Subject Classification:* 53C15.
*Key Words and Phrases:* para-hyperhermitian structure, tangent bundle, mixed 3-structure, semi-Riemannian submersion.

## 1. Introduction

The para-hyperhermitian structures arise in a natural way in theoretical physics, both in string theory and integrable systems ([10], [12], [13], [15], [25]). This kind of structures have been intensively studied with different names by many authors (see [2], [14], [18], [19], [22], [27] and more).

In this paper we construct some classes of manifolds endowed with almost para-hyperhermitian structures. The paper is organized as follows. In section 2 we recall the definition and fundamental properties of almost para-hyperhermitian manifolds.

In Section 3 we give an almost para-hyperhermitian structure on the tangent bundle of an almost para-hermitian manifold and study its integrability.

The concept of mixed 3-structure has been introduced in [17]. In section 4 we study the manifolds endowed with such structures and prove that $M$ is a manifold with a mixed 3-structure, then $\overline{M}=M\times\mathbb{R}$ can be endowed with an almost para-hyperhermitian structure. We construct also an almost para-hyperhermitian structures on a principal circle bundle over a smooth manifold endowed with a metric mixed 3-structure.

## 2. Preliminaries on almost para-hyperhermitian manifolds

An almost product structure on a smooth manifold $M$ is a tensor field $P$ of type (1,1) on $M$, $P\neq\pm Id$, such that:

$$P^2=Id. \tag{1}$$

An almost complex structure on a smooth manifold $M$ is a tensor field $J$ of type (1,1) on $M$ such that:

$$J^2=-Id. \tag{2}$$

The authors are partially supported by grant D11-22 CEEX 2006-2008. This paper has been presented in the "4th German-Romanian Seminar on Geometry" Dortmund, Germany, 15-18 July 2007.

An almost para-hypercomplex structure on a smooth manifold $M$ is a triple $H = (J_\alpha)_{\alpha=\overline{1,3}}$, where $J_1$ is an almost complex structure on $M$ and $J_2$, $J_3$ are almost product structures on $M$, satisfying:

$$J_2 J_1 = -J_1 J_2 = J_3. \tag{3}$$

In this case $(M, H)$ is said to be an almost para-hypercomplex manifold.

It is easy to see that any almost para-hypercomplex manifold is of dimension $4n$, for a non-negative integer $n$.

A semi-Riemannian metric $g$ on $(M, H)$ is said to be para-hyperhermitian if it satisfies:

$$g(J_\alpha X, J_\alpha Y) = \epsilon_\alpha g(X, Y), \forall \alpha = \overline{1,3} \tag{4}$$

for all vector fields $X$,$Y$ on $M$, where $\epsilon_1 = 1, \epsilon_2 = \epsilon_3 = -1$.

Moreover, $(M, g, H)$ is said to be an almost para-hyperhermitian manifold. It is clear that the signature of $g$ is $(2n, 2n)$ because, at each point of $M$, there is a local pseudo-orthonormal frame field, called adapted frame, given in the following way:

$$\{E_1, ..., E_m, J_1E_1, ..., J_1E_m, J_2E_1, ..., J_2E_m, J_3E_1, ..., J_3E_m\}.$$

If $h$ is an arbitrary semi-Riemannian metric on an almost para-hypercomplex manifold $(M, H)$, then we can define always a para-hyperhermitian metric $g$ on $M$ by:

$$g(X, Y) = \frac{1}{4}[h(X, Y) + \sum_{\alpha=1}^{3} \epsilon_\alpha h(J_\alpha X, J_\alpha Y)] \tag{5}$$

for all vector fields $X$,$Y$ on $M$.

An almost para-hypercomplex manifold $(M, H)$ is said to be a para-hypercomplex manifold if each $J_\alpha$, $\alpha = 1, 2, 3$, is integrable, that is, if the corresponding Nijenhuis tensors:

$$N_\alpha(X, Y) = [J_\alpha X, J_\alpha Y] - J_\alpha[X, J_\alpha Y] - J_\alpha[J_\alpha X, Y] - \epsilon_\alpha[X, Y] \tag{6}$$

$\alpha = 1, 2, 3$, vanish for all vector fields $X$,$Y$ on $M$. In this case $H$ is said to be a para-hypercomplex structure on $M$.

We remark that if two of the structures $J_1, J_2, J_3$ are integrable, then the third structure is also integrable because we have:

$$\begin{aligned} 2N_\alpha(X, Y) &= N_\beta(J_\gamma X, J_\gamma Y) + N_\gamma(J_\beta X, J_\beta Y) - J_\beta N_\gamma(J_\beta X, Y) - J_\beta N_\gamma(X, J_\beta Y) \\ &\quad -J_\gamma N_\beta(J_\gamma X, Y) - J_\gamma N_\beta(X, J_\gamma Y) + \epsilon_\alpha\epsilon_\beta N_\beta(X, Y) + \epsilon_\alpha\epsilon_\gamma N_\gamma(X, Y) \end{aligned} \tag{7}$$

for any even permutation $(\alpha, \beta, \gamma)$ of (1,2,3), where $\epsilon_1 = 1, \epsilon_2 = \epsilon_3 = -1$.

An almost hermitian paracuaternionic manifold is a triple $(M, \sigma, g)$, where $M$ is a smooth manifold, $\sigma$ is a rank 3-subbundle of $End(TM)$ which is locally spanned by an almost para-hypercomplex structure $H = (J_\alpha)_{\alpha=\overline{1,3}}$ and $g$ is a para-hyperhermitian metric in respect with $H$.

## 3. An almost para-hyperhermitian structure on the tangent bundle of an almost para-hermitian manifold

An almost para-hermitian structure on a differentiable manifold $M$ is a pair $(P, g)$, where $P$ is an almost product structure on $M$ and $g$ is a semi-Riemannian metric on $M$ satisfying:

$$g(PX, PY) = -g(X, Y), \tag{8}$$

for all vector fields $X$,$Y$ on $M$.

In this case, $(M, P, g)$ is said to be an almost para-hermitian manifold. It is easy to see that the dimension of $M$ is even. Moreover, if $\nabla P = 0$ then $(M, P, g)$ is said to be a para-Kähler manifold.

We remark that this kind of manifolds appeared for the first time in [22].

Now, let $(M, P, g)$ be an almost para-hermitian manifold and $TM$ be the tangent bundle, endowed with the Sasakian metric:

$$G(X, Y) = (g(KX, KY) + g(\pi_* X, \pi_* Y)) \circ \pi$$

for all vector fields $X, Y$ on $TM$, where $\pi$ is the natural projection of $TM$ onto $M$ and $K$ is the connection map (see [11]).

We remark that if $X \in \Gamma(TM)$, then there exists exactly one vector field on $TM$ called the "horizontal lift" (resp. "vertical lift") of $X$ such that for all $t \in TM$:

$$\pi_* X_t^h = X_{\pi(t)},\ \pi_* X_t^v = 0_{\pi(t)},\ KX_t^h = 0_{\pi(t)},\ KX_t^v = X_{\pi(t)}.$$

*Remark* 3.1. It is immediately that for any affine connection $\nabla$ on $M$ we have:

$$[X^h, Y^h] = [X, Y]^h - (R(X, Y)t)^v,\ [X^v, Y^v] = 0,$$
$$[X^h, Y^v] = (\nabla_X Y)^v,\ [X^v, Y^h] = -(\nabla_Y X)^v,$$

for all $X, Y \in T_t(TM)$, $t \in TM$, where $R$ is the curvature tensor of $\nabla$ on $M$.

**Theorem 3.2.** *Let $(M, P, g)$ be an almost para-hermitian manifold. Then $TM$ admits an almost para-hypercomplex structure $H$ which is para-hyperhermitian in respect to $G$.*

*Proof.* We define three tensor fields $J_1, J_2, J_3$ on $TM$ by the equalities:

$$J_1 X^h = X^v, J_1 X^v = -X^h, \tag{9}$$

$$J_2 X^h = (PX)^v, J_2 X^v = (PX)^h, \tag{10}$$

$$J_3 X^h = (PX)^h, J_3 X^v = -(PX)^v. \tag{11}$$

We can easily see that we have:

$$J_1^2 = -J_2^2 = -J_3^2 = -Id,$$
$$J_2 J_1 = -J_1 J_2 = J_3$$

and

$$G(J_1 X, J_1 Y) = -G(J_2 X, J_2 Y) = -G(J_3 X, J_3 Y) = G(X, Y).$$

□

**Corollary 3.3.** *The tangent bundle of the pseudosphere $S_3^6$ admits an almost para-hyperhermitian structure.*

*Proof.* The assertion is clear from above Theorem because the pseudosphere $S_3^6$ can be endowed with an almost para-hermitian structure (see [3]). □

**Corollary 3.4.** *Let $(M, g)$ be a semi-Riemannian manifold and $T^*M$ its cotangent bundle. Then the tangent bundle of the cotangent bundle $TT^*M$ admits an almost para-hyperhermitian structure.*

*Proof.* The assertion is clear from Theorem 3.2 because the cotangent bundle $T^*M$ can be endowed with an almost para-hermitian structure (see [26]). □

**Theorem 3.5.** *Let $(M,P,g)$ be an almost para-hermitian manifold. Then the almost para-hypercomplex structure $H=(J_\alpha)_{\alpha=\overline{1,3}}$ on $TM$ given by Theorem 3.2 is integrable if and only if $(M,P)$ is a flat para-Kähler manifold.*

*Proof.* Using Remark 3.1 we deduce that the Nijenhuis tensor of $J_1$ is given by:

$$\begin{aligned} N_1(X^h,Y^h) &= [J_1X^h,J_1Y^h]-J_1[X^h,J_1Y^h]-J_1[J_1X^h,Y^h]-[X^h,Y^h] \\ &= J_1(\nabla_YX)^v-J_1(\nabla_XY)^v-[X,Y]^h+(R(X,Y)t)^v \\ &= (R(X,Y)t)^v, \end{aligned} \tag{12}$$

$$\begin{aligned} N_1(X^v,Y^v) &= [J_1X^v,J_1Y^v]-J_1[X^v,J_1Y^v]-J_1[J_1X^v,Y^v]-[X^v,Y^v] \\ &= [X,Y]^h-(R(X,Y)t)^v-J_1(\nabla_YX)^v+J_1(\nabla_XY)^v \\ &= -(R(X,Y)t)^v, \end{aligned} \tag{13}$$

$$\begin{aligned} N_1(X^h,Y^v) &= [J_1X^h,J_1Y^v]-J_1[X^h,J_1Y^v]-J_1[J_1X^h,Y^v]-[X^h,Y^v] \\ &= (\nabla_YX)^v+J_1([X,Y]^h-(R(X,Y)t)^v)-(\nabla_XY)^v \\ &= (R(X,Y)t)^h, \end{aligned} \tag{14}$$

$$\begin{aligned} N_1(X^v,Y^h) &= [J_1X^v,J_1Y^h]-J_1[X^v,J_1Y^h]-J_1[J_1X^v,Y^h]-[X^v,Y^h] \\ &= -(\nabla_XY)^v+J_1([X,Y]^h-(R(X,Y)t)^v)+(\nabla_YX)^v \\ &= (R(X,Y)t)^h. \end{aligned} \tag{15}$$

for all $X,Y\in T_t(TM)$, $t\in TM$.

Similarly as above, we deduce:

$$N_2(X^h,Y^h)=(P(\nabla_YP)X-P(\nabla_XP)Y)^h-(R(X,Y)t)^v, \tag{16}$$

$$N_2(X^v,Y^v)=((\nabla_{PX}P)Y-(\nabla_{PY}P)X)^h-(R(PX,PY)t)^v, \tag{17}$$

$$N_2(X^h,Y^v)=-(P(\nabla_XP)Y+(\nabla_{PY}P)X)^v+(PR(X,PY)t)^h, \tag{18}$$

$$N_2(X^v,Y^h)=((\nabla_{PX}P)Y+P(\nabla_YP)X)^v+(PR(PX,Y)t)^h \tag{19}$$

and

$$\begin{aligned} N_3(X^h,Y^h) &= ((\nabla_XP)PY-(\nabla_{PY}P)X+(\nabla_{PX}P)Y+P(\nabla_YP)X)^h \\ &\quad -(R(X,Y)t+R(PX,PY)t+PR(PX,Y)t+PR(X,PY)t)^v, \end{aligned} \tag{20}$$

$$N_3(X^v,Y^v)=0, \tag{21}$$

$$N_3(X^h,Y^v)=(-(\nabla_{PX}P)Y+(\nabla_XP)(PY))^v, \tag{22}$$

$$N_3(X^v,Y^h)=((\nabla_{PY}P)X-(\nabla_YP)(PX))^v. \tag{23}$$

The proof is now complete from (12)-(23). □

**Corollary 3.6.** *Let $(M,P,g)$ be an almost para-hermitian manifold and $H=(J_\alpha)_{\alpha=\overline{1,3}}$ the almost para-hypercomplex structure on $TM$ given in Theorem 3.2. If $J_2$ is integrable, then $H$ is a para-hypercomplex structure on $TM$.*

*Proof.* The proof is clear from Theorem 3.5. □

## 4. Almost para-hyperhermitian structures and manifolds endowed with mixed 3-structures

**Definition 4.1.** Let $M$ be a differentiable manifold equipped with a triple $(\phi, \xi, \eta)$, where $\phi$ is a a field of endomorphisms of the tangent spaces, $\xi$ is a vector field and $\eta$ is a 1-form on $M$. If we have:

$$\phi^2 = -\epsilon I + \eta \otimes \xi, \ \eta(\xi) = \epsilon \tag{24}$$

then we say that:
i. $(\phi, \xi, \eta)$ is an almost contact structure on $M$, if $\epsilon = 1$ (cf. [5]).
ii.$(\phi, \xi, \eta)$ is a Lorentzian almost paracontact structure on $M$, if $\epsilon = -1$ (cf. [24]).

We remark that many authors also include in the above definition the conditions that $\phi\xi = 0$ and $\eta \circ \phi = 0$, although these are deducible from the conditions (24) (see [5]).

**Definition 4.2.** ([17]) Let $M$ be a differentiable manifold which admits an almost contact structure $(\phi_1, \xi_1, \eta_1)$ and two Lorentzian almost paracontact structures $(\phi_2, \xi_2, \eta_2)$ and $(\phi_3, \xi_3, \eta_3)$, satisfying the following conditions:
i.

$$\eta_\alpha(\xi_\beta) = 0, \forall \alpha \neq \beta; \tag{25}$$

ii.

$$\phi_\alpha(\xi_\beta) = -\phi_\beta(\xi_\alpha) = \epsilon_\gamma \xi_\gamma; \tag{26}$$

iii.

$$\eta_\alpha \circ \phi_\beta = -\eta_\beta \circ \phi_\alpha = \epsilon_\gamma \eta_\gamma; \tag{27}$$

iv.

$$\phi_\alpha \phi_\beta - \eta_\beta \otimes \xi_\alpha = -\phi_\beta \phi_\alpha + \eta_\alpha \otimes \xi_\beta = \epsilon_\gamma \phi_\gamma, \tag{28}$$

where in (26), (27) and (28), $(\alpha, \beta, \gamma)$ is an even permutation of (1,2,3) and $\epsilon_1 = 1, \epsilon_2 = \epsilon_3 = -1$.

Then the manifold $M$ is said to have a mixed 3-structure $(\phi_\alpha, \xi_\alpha, \eta_\alpha)_{\alpha=\overline{1,3}}$.

**Definition 4.3.** If a manifold $M$ with a mixed 3-structure $(\phi_\alpha, \xi_\alpha, \eta_\alpha)_{\alpha=\overline{1,3}}$ admits a semi-Riemannian metric $g$ such that:

$$g(\phi_\alpha X, \phi_\alpha Y) = \epsilon_\alpha g(X, Y) - \eta_\alpha(X)\eta_\alpha(Y), \tag{29}$$

and

$$g(X, \xi_\alpha) = \eta_\alpha(X) \tag{30}$$

for all $X, Y \in \Gamma(TM)$ and $\alpha = 1, 2, 3$, then we say that $M$ has a metric mixed 3-structure and $g$ is called a compatible metric.

Moreover, if $(\phi_1, \xi_1, \eta_1, g)$ is a Sasakian structure, i.e.(see [5]):

$$(\nabla_X \phi_1)Y = g(X, Y)\xi_1 - \eta_1(Y)X$$

and $(\phi_2, \xi_2, \eta_2, g)$, $(\phi_3, \xi_3, \eta_3, g)$ are Lorentzian para-Sasakian structures, i.e.(see [24]):

$$(\nabla_X \phi_2)Y = g(\phi_2 X, \phi_2 Y)\xi_2 + \eta_2(Y)\phi_2^2 X,$$
$$(\nabla_X \phi_3)Y = g(\phi_3 X, \phi_3 Y)\xi_3 + \eta_3(Y)\phi_3^2 X,$$

where $\nabla$ is the Levi-Civita connection of $g$, then $((\phi_\alpha, \xi_\alpha, \eta_\alpha)_{\alpha=\overline{1,3}}, g)$ is said to be a mixed Sasakian 3-structure on $M$.

**Proposition 4.4.** *Any manifold $M$ with a mixed 3-structure $(\phi_\alpha, \xi_\alpha, \eta_\alpha)_{\alpha=\overline{1,3}}$ admits a compatible semi-Riemannian metric.*

*Proof.* A such type of metric can be constructed from any semi-Riemannian metric $f$ on $M$, in four step:

i. First we define the metric $u$ by:

$$u(X,Y) = f(\phi_1^2 X, \phi_1^2 Y) + \eta_1(X)\eta_1(Y)$$

and we can see that $u(X, \xi_1) = \eta_1(X)$.

ii. We define now a new metric $v$ by:

$$v(X,Y) = u(\phi_2^2 X, \phi_2^2 Y) - \eta_2(X)\eta_2(Y)$$

and we have that $v(X, \xi_\alpha) = \eta_\alpha(X)$, for $\alpha = 1, 2$.

iii. We define a new metric $h$ by:

$$h(X,Y) = v(\phi_3^2 X, \phi_3^2 Y) - \eta_3(X)\eta_3(Y)$$

and we can easily see that $h(X, \xi_\alpha) = \eta_\alpha(X)$, for $\alpha = 1, 2, 3$.

iv. Finally we define the metric $g$ by:

$$g(X,Y) = \frac{1}{4}[h(X,Y) + \sum_{\alpha=1}^{3} \epsilon_\alpha [h(\phi_\alpha X, \phi_\alpha Y) + \eta_\alpha(X)\eta_\alpha(Y)]$$

and a straightforward verification shows that $g$ provides a compatible semi-Riemannian metric on $(M, (\phi_\alpha, \xi_\alpha, \eta_\alpha)_{\alpha=\overline{1,3}})$. □

*Remark* 4.5. If $(M^{4n+3}, (\phi_\alpha, \xi_\alpha, \eta_\alpha)_{\alpha=\overline{1,3}}, g)$ is a manifold with a metric mixed 3-structure, then it is easy to see that the signature of $g$ is $(2n+1, 2n+2)$ because one can check that, at each point of $M$, there always exists a pseudo-orthonormal frame field given in the following way:

$$\{(E_i, \phi_1 E_i, \phi_2 E_i, \phi_3 E_i)_{i=1,n}, \xi_1, \xi_2, \xi_3\}.$$

*Example* 4.6. 1. It is easy to see that if we define $(\phi_\alpha, \xi_\alpha, \eta_\alpha)_{\alpha=\overline{1,3}}$ in $\mathbb{R}^3$ by their matrices:

$$\phi_1 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix},\ \phi_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix},\ \phi_3 = \begin{pmatrix} 0 & -1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

$$\xi_1 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix},\ \xi_2 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix},\ \xi_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix},$$

$$\eta_1 = \begin{pmatrix} 0 & 1 & 0 \end{pmatrix},\ \eta_2 = \begin{pmatrix} -1 & 0 & 0 \end{pmatrix},\ \eta_3 = \begin{pmatrix} 0 & 0 & -1 \end{pmatrix},$$

then $(\phi_\alpha, \xi_\alpha, \eta_\alpha)_{\alpha=\overline{1,3}}$ is a mixed 3-structure on $\mathbb{R}^3$.

We define now $(\phi'_\alpha, \xi'_\alpha, \eta'_\alpha)_{\alpha=\overline{1,3}}$ in $\mathbb{R}^{4n+3}$ by:

$$\phi'_\alpha = \begin{pmatrix} \phi_\alpha & 0 \\ 0 & J_\alpha \end{pmatrix}, \xi'_\alpha = \begin{pmatrix} \xi_\alpha \\ 0 \end{pmatrix}, \eta'_\alpha = \begin{pmatrix} \eta_\alpha & 0 \end{pmatrix},$$

for $\alpha = 1, 2, 3$, where $J_1$ is the almost complex structure on $\mathbb{R}^{4n}$ given by:

$$J_1((x_i)_{i=\overline{1,4n}}) = (-x_2, x_1, -x_4, x_3, ..., -x_{4n-2}, x_{4n-3}, -x_{4n}, x_{4n-1}), \tag{31}$$

and $J_2,\ J_3$ are almost product structures on $\mathbb{R}^{4n}$ defined by:

$$J_2((x_i)_{i=\overline{1,4n}}) = (-x_{4n-1}, x_{4n}, -x_{4n-3}, x_{4n-2}, ..., -x_3, x_4, -x_1, x_2), \tag{32}$$

$$J_3((x_i)_{i=\overline{1,4n}}) = (x_{4n}, x_{4n-1}, x_{4n-2}, x_{4n-3}, ..., x_4, x_3, x_2, x_1). \tag{33}$$

Since $J_2J_1 = -J_1J_2 = J_3$, it is easily checked that $(\phi'_\alpha, \xi'_\alpha, \eta'_\alpha)_{\alpha=\overline{1,3}}$ is a mixed 3-structure on $\mathbb{R}^{4n+3}$.

2. Let $(\overline{M}, \overline{g})$ be a $(m+2)$-dimensional semi-Riemannian manifold with index $q \in \{1, 2, \ldots, m+1\}$ and let $(M, g)$ be a hypersurface of $\overline{M}$, with $g = \overline{g}_{|M}$. We say that $M$ is a lightlike hypersurface of $\overline{M}$ if $g$ is of constant rank $m$ (see [4]).

We consider the vector bundle $TM^\perp$ whose fibres are defined by:

$$T_pM^\perp = \{Y_p \in T_p\overline{M} | \overline{g}_p(X_p, Y_p) = 0, \forall X_p \in T_pM\}, \forall p \in M.$$

If $S(TM)$ is the complementary distribution of $TM^\perp$ in $TM$, which is called the screen distribution, then there exists a unique vector bundle $ltr(TM)$ of rank 1 over $M$ so that for any non-zero section $\xi$ of $TM^\perp$ on a coordinate neighborhood $U \subset M$, there exists a unique section $N$ of $ltr(TM)$ on $U$ satisfying:

$$\overline{g}(N, \xi) = 1,\ \overline{g}(N, N) = \overline{g}(W, W) = 0,\ \forall W \in \Gamma(S(TM)_{|U})$$

(see [4]).

In a lightlike hypersurface $M$ of an almost hermitian paraquaternionic manifold $(\overline{M}, \overline{g}, \sigma)$ such that $\xi$ and $N$ are globally defined on $M$, there is a mixed 3-structure (see [17]).

3. The unit sphere $S^{4n+3}_{2n+1}$ is the canonical example of manifold with a mixed Sasakian 3-structure. This structure is obtained by taking $S^{4n+3}_{2n+1}$ as hypersurface of $(\mathbb{R}^{4n+4}_{2n+2}, \overline{g})$. It is easy to see that on the tangent spaces $T_pS^{4n+3}_{2n+1}$, $p \in S^{4n+3}_{2n+1}$, the induced metric $g$ is of signature $(2n+1, 2n+2)$.

If $(J_\alpha)_{\alpha=\overline{1,3}}$ is the canonical paraquaternionic structure on the $\mathbb{R}^{4n+4}_{2n+2}$ and $N$ is the unit normal vector field to the sphere, we can define three vector fields on $S^{4n+3}_{2n+1}$ by:

$$\xi_\alpha = -J_\alpha N,\ \alpha = 1, 2, 3.$$

If $X$ is a tangent vector to the sphere then $J_\alpha X$ uniquely decomposes onto the part tangent to the sphere and the part parallel to $N$. Denote this decomposition by:

$$J_\alpha X = \phi_\alpha X + \eta_\alpha(X)N.$$

This defines the 1-forms $\eta_\alpha$ and the tensor fields $\phi_\alpha$ on $S^{4n+3}_{2n+1}$, where $\alpha = 1, 2, 3$.

Now we can easily see that that $(\phi_\alpha, \xi_\alpha, \eta_\alpha)_{\alpha=\overline{1,3}}$ is a mixed Sasakian 3-structure on $S^{4n+3}_{2n+1}$.

4. Since we can recognize the unit sphere $S^{4n+3}_{2n+1}$ as the projective space $P^{4n+3}_{2n+1}(\mathbb{R})$, by identifying antipodal points, we have also that $P^{4n+3}_{2n+1}(\mathbb{R})$ admits a mixed Sasakian 3-structure.

**Theorem 4.7.** *If $M$ is a manifold with a mixed 3-structure $(\phi_\alpha, \xi_\alpha, \eta_\alpha)_{\alpha=\overline{1,3}}$ then $\overline{M} = M \times \mathbb{I}$ can be endowed with an almost para-hyperhermitian structure, where $\mathbb{I}$ is $\mathbb{R}$ or some open interval in $\mathbb{R}$.*

*Proof.* We define three tensor fields $J_1, J_2, J_3$ on $\overline{M}$ by the equalities:

$$J_\alpha = \begin{pmatrix} \phi_\alpha & \frac{\xi_\alpha}{f} \\ -f\eta_\alpha & 0 \end{pmatrix}, \alpha = 1, 2, 3. \tag{34}$$

where $f$ is a positive function on $\mathbb{I}$.

By a straightforward computation one can check:

$$J_1^2 = -J_2^2 = -J_3^2 = -Id,$$
$$J_2 J_1 = -J_1 J_2 = J_3$$

and the proof is complete because we can construct now a para-hyperhermitian metric on $\overline{M} = M \times \mathbb{R}$ from any arbitrary semi-Riemannian metric. □

**Corollary 4.8.** *$S^{4n+3}_{2n+1} \times \mathbb{I}$ and $P^{4n+3}_{2n+1}(\mathbb{R}) \times \mathbb{I}$ can be endowed with almost para-hyperhermitian structures, where $\mathbb{I}$ is $\mathbb{R}$ or some open interval in $\mathbb{R}$.*

*Proof.* From Example 4.6 we have that $S^{4n+3}_{2n+1}$ and $P^{4n+3}_{2n+1}(\mathbb{R})$ are mixed 3-Sasakian manifolds and, consequently, each structure $J_\alpha$ given by (34) is integrable. □

**Corollary 4.9.** *Let $M$ be a manifold endowed with a mixed 3-structure $(\phi_\alpha, \xi_\alpha, \eta_\alpha)_{\alpha=\overline{1,3}}$. Then the dimension of $M$ is $4n+3$, where $n$ is a non-negative integer.*

*Proof.* The assertion is clear from Theorem 4.7. □

*Remark* 4.10. If $M^{4n+3}$ is a manifold endowed with a mixed 3-Sasakian structure $((\phi_\alpha, \xi_\alpha, \eta_\alpha)_{\alpha=\overline{1,3}}, g)$, then we can define a para-hyper-Kähler structure $\{J_\alpha\}_{\alpha=\overline{1,3}}$ on the cone $(C(M), \overline{g}) = (M \times \mathbb{R}_+, dr^2 + r^2 g)$, by:

$$\left\{ \begin{array}{rcl} J_\alpha X & = & \phi_\alpha X - \eta_\alpha(X)\Phi \\ J_\alpha \Phi & = & \xi_\alpha \end{array} \right. \tag{35}$$

for any vector field $X \in \Gamma(TM)$ and $\alpha = 1, 2, 3$, where $\Phi = r\partial_r$ is the Euler field on $C(M)$.

Moreover, conversely, if a cone $(C(M), \overline{g}) = (M \times \mathbb{R}_+, dr^2 + r^2 g)$ admits a para-hyper-Kähler structure $\{J_\alpha\}_{\alpha=\overline{1,3}}$, then we can identify $M$ with $M \times \{1\}$ and we have a mixed 3-Sasakian structure $((\phi_\alpha, \xi_\alpha, \eta_\alpha)_{\alpha=\overline{1,3}}, g)$ on $M$ given by:

$$\xi_\alpha = J_\alpha(\partial_r),\ \phi_\alpha X = \nabla_X \xi_\alpha,\ \eta_\alpha(X) = g(\xi_\alpha, X), \tag{36}$$

for any vector field $X \in \Gamma(TM)$ and $\alpha = 1, 2, 3$.

Finally, since a para-hyper-Kähler manifold is Ricci-flat, we conclude that $M$ is an Einstein space with Einstein constant $\lambda = 4n + 2$ (see [7]).

*Remark* 4.11. Let $P = P(M, \pi, S^1)$ be a principal circle bundle over a smooth manifold $M^{4n+3}$ with a metric mixed 3-structure $((\phi_\alpha, \xi_\alpha, \eta_\alpha)_{\alpha=\overline{1,3}}, g)$.

Let $g'$ be a semi-Riemannian metric on $P$ such that $\pi : (P, g') \to (M, g)$ is a semi-Riemannian submersion. Putting $\mathcal{V}_x = Ker\ \pi_{*x}$, for any $x \in M$, we obtain an integrable distribution $\mathcal{V}$, which is called vertical distribution and corresponds to the foliation of $M$ determined by the fibres of $\pi$. The complementary distribution $\mathcal{H}$ of $\mathcal{V}$, determined by the semi-Riemannian metric $g'$, is called horizontal distribution. We have now the decomposition:

$$TP = \mathcal{H} \oplus \mathcal{V}.$$

We recall that the sections of $\mathcal{V}$, respectively $\mathcal{H}$, are called the vertical vector fields, respectively horizontal vector fields. An horizontal vector field $X^h$ on $P$ is said to be basic if $X^h$ is $\pi$-related to a vector field $X'$ on $M$. It is clearly that every vector field $X$ on $M$ has a unique horizontal lift $X^h$ to $P$ and $X^h$ is basic.

**Theorem 4.12.** *There is an almost para-hyperhermitian structure on any principal circle bundle $P = P(M, \pi, S^1)$ over a manifold $M$ with a metric mixed 3-structure $((\phi_\alpha, \xi_\alpha, \eta_\alpha)_{\alpha=\overline{1,3}}, g)$.*

*Proof.* Making use of the metric mixed 3-structure on $M$ we can define three tensor fields $J_1, J_2, J_3$ on $P$ by the equalities:

$$\left\{ \begin{array}{rcl} J_\alpha X^h & = & (\phi_\alpha X)^h + \eta_\alpha(X)\Theta \\ J_\alpha \Theta & = & -(\xi_\alpha)^h \end{array} \right. \tag{37}$$

for $\alpha = 1, 2, 3$, where $\Theta$ is the nowhere vanishing vertical vector field on $P$ which generates the $S^1$ action on $P$.

Now, we can easily see that we have:

$$J_1^2 = -J_2^2 = -J_3^2 = -Id,$$

$$J_2 J_1 = -J_1 J_2 = J_3$$

and the proof is now complete since we can construct a para-hyperhermitian metric from any arbitrary semi-Riemannian metric on $P$. □

**Corollary 4.13.** *$\mathbb{R}^{4n+3}_{2n+1} \times S^1$, $S^{4n+3}_{2n+1} \times S^1$ and $P^{4n+3}_{2n+1}(\mathbb{R}) \times S^1$ can be endowed with almost para-hyperhermitian structures.*

*Proof.* The assertion is clear from Example 4.6 and Theorem 4.12. □

## References


[1] D.V. Alekseevsky, N. Blazic, V. Cortés, S. Vukmirovic, *A class of Osserman spaces*, J. Geom. Phys. 53, No. 3 (2005), 345-353.

[2] D. Alekseevsky, Y. Kamishima, *Quaternionic and para-quaternionic CR structure on (4n+3)-dimensional manifolds*, Central European J. Math. 2(5) (2004), 732-753.

[3] C.L. Bejan, *Some examples of manifolds with hyperbolic structures*, Rend. Mat. Appl., VII. Ser. 14, No.4 (1994), 557-565.

[4] A. Bejancu, K.L. Duggal, *Lightlike submanifolds of semi-Riemannian manifolds and its application*, Kluwer, Dortrecht 1996.

[5] D.E. Blair, *Contact manifolds in Riemannian Geometry*, Lectures Notes in Math. 509, Springer-Verlag, 1976.

[6] N. Blazic, S. Vukmirovic, *Para-hypercomplex structures on a four-dimensional Lie group*, Contemporary geometry and related topics. Proceedings of the workshop, Belgrade, Yugoslavia, May 15-21, 2002. River Edge, NJ: World Scientific. 41-56 (2004).

[7] C. Boyer, K. Galicki, *3-Sasakian manifolds*, Surveys in differential geometry: essays on Einstein manifolds. Lectures on geometry and topology, Cambridge, MA: International Press. Surv. Differ. Geom., Suppl. J. Differ. Geom. 6, 123-184 (1999).

[8] C. Boyer, K. Galicki, B. Mann, *Hypercomplex structures on Stiefel manifolds*, Ann. Global Anal. Geom. 14, No.1 (1996), 81-105.

[9] F.M Cabrera, *Almost hyper-Hermitian structures in bundle spaces over manifolds with almost contact 3-structure*, Czech. Math. J. 48 (1998), No.3, 545-563.

[10] V. Cortés, C. Mayer, T. Mohaupt, F. Saueressig *Special geometry of euclidean supersymmetry II. Hypermultiplets and the c-map*, J. High Energy Phys. 06 (2005) 025, hep-th/0503094.

[11] P. Dombrowski, *On the geometry of the tangent bundle*, J. Reine und Angew. Math., 210 (1962), 73-88.

[12] M. Dunajski, *Hyper-complex four-manifolds from Tzitzeica equation*, J. Math. Phys. 43 (2002), 651-658.

[13] M. Dunajski, S. West *Anti-self-dual conformal structures in neutral signature*, e-print math.DG/0610280 (2006).

[14] N. Hitchin, *Hypersymplectic quotients*, Acta Acad. Sci. Tauriensis 124 supl., (1990), 169-180.

[15] C. M. Hull, *Actions for (2,1) Sigma Models and Strings*, Nucl. Phys. B509 (1998) 252-272.

[16] S. Ianuş, *Sulle strutture canoniche dello spazio fibrato tangente di una varieta riemanniana*, Rend. Mat., 6 (1973), 75-96.

[17] S. Ianuş, R. Mazzocco, G.E. Vîlcu, *Real lightlike hypersurfaces of paraquaternionic Kähler manifolds"*, Mediterr. J. Math., 3 (2006), 581-592.

[18] S. Ivanov, S. Zamkovoy, *Para-hermitian and para-quaternionic manifolds.*, Differential Geom. Appl., 23 (2005), 205-234.

[19] H. Kamada, *Neutral hyper-Kähler structures on primary Kodaira surfaces*, Tsukuba J. Math. 23 (1999), 321-332.
[20] O. Kowalski, *Curvature of the induced Riemannian metric on the tangent bundle of a Riemannian manifold*, J. Reine Angew. Math. 250 (1971), 124-129.
[21] Y.Y. Kuo, *On almost contact 3-structure*, Tohoku Math. J., II. Ser. 22 (1970), 325-332.
[22] P. Libermann, *Sur le problème d'équivalence de certaines structures infinitésimales*, Ann. Mat. Purra Appl., 36 (1954), 27-120.
[23] M. Manev, *Tangent bundles with Sasaki metric and almost hypercomplex pseudo-Hermitian structure*, Topics in almost Hermitian geometry and related fields. Proceedings in honor of Prof. K. Sekigawa's 60th birthday, Niigata, Japan, November 1-3, 2004. Hackensack, NJ: World Scientific. 170-185 (2005).
[24] K. Matsumoto, *On Lorentzian paracontact manifolds*, Bull. Of Yamagata Univ. Nat. Sci., 12(2) (1989), 151-156.
[25] H. Ooguri, C. Vafa, *Geometry of N=2 strings*, Nucl. Phys. B 361 (1991), 469-518.
[26] V. Oproiu, N. Papaghiuc, G. Mitric, *Some classes of parahermitian structures on cotangent bundles*, An. Stiint. Univ. Al. I. Cuza Iasi, Ser. Noua, Mat. 43, No.1 (1996), 7-22.
[27] S. Vukmirović, *Para-quaternionic reduction*, e-print math. DG/03044424 (2003).

Stere Ianuş
*University of Bucharest,*
*Department of Mathematics,*
*C.P. 10-119, Post. Of. 10, Bucharest 72200, România*
e-mail: ianus@gta.math.unibuc.ro

Gabriel Eduard Vîlcu
*"Petroleum-Gas" University of Ploieşti,*
*Department of Mathematics and Computer Science,*
*Bulevardul Bucureşti, Nr. 39, Ploieşti, România*
e-mail: gvilcu@mail.upg-ploiesti.ro